

NEW

Extinguish™ XL

Arylex™ active

HERBICIDE

Delivering a combination of high performance and value, Extinguish™ XL herbicide with Arylex™ active is a powerful new solution to your weed control problems.

WHY USE EXTINGUISH™ XL?

- **Reliable, economical control** of key broadleaf weeds like cleavers, dandelion, round-leaved mallow, stork's-bill and narrow-leaved hawk's beard.
- Convenient all-in-one liquid formulation and compact case size.
- Extinguish XL delivers **consistent broadleaf weed control**.
- **Wide window** of application provides control of small and large weeds and excellent crop safety from the 2-leaf to just prior to flag leaf stage.
- **Multiple modes of action** control key broadleaf weeds to delay the onset of herbicide resistance.

**BROADLEAF
WEEDS
CONTROLLED**

- Annual Sow-thistle¹
- Canada
- Fleabane²
- Chickweed^{1,3}
- Cleavers^{1,3}
- Common ragweed²
- Cow Cockle¹
- Dandelion
- Flixweed
- Lamb's quarters³
- Narrow-leaved hawk's beard
- Redroot pigweed¹

- Round-leaved mallow
- Shepherd's purse¹
- Smartweed
- Stinkweed¹
- Stork's-bill
- Velvetleaf
- Volunteer alfalfa
- Volunteer canola⁴
- Volunteer flax
- Wild Buckwheat¹
- Wild mustard¹

**GRASS WEEDS
CONTROLLED**

- Barnyard grass

**WEEDS
SUPPRESSED**

- Hemp-nettle^{1,3}
- Kochia²
- Night flowering catchfly
- Scentless chamomile
- Perennial sow-thistle
- Canada thistle
- White cockle

¹ Weed controlled by multiple effective modes of action (except for Group 2 resistant biotypes)

² Including Group 2 and 9 resistant biotypes

³ Including Group 2 resistant biotypes

⁴ Including all herbicide-resistant biotypes

APPLICATION GUIDELINES

CROPS Barley Durum wheat Spring wheat Winter wheat	PACKAGING	Case: 2 x 6.74 L jugs Drum: 80.9 L
	RATES	337 mL/ac
	ACRES TREATED	20 ac/jug 240 ac/drum
	WATER VOLUME	• Ground: 20-40L/ac (5-10 US gal/ac)
	TIMING	Crop stage: 2-leaf to just prior to flag leaf emergence Weed stage: 1-leaf to 8-leaf
	RAINFEST	1 hour
	TANK-MIX HERBICIDES	• Axial® • Everest® 3.0 • Simplicity™ ⁵ herbicide • Simplicity™ GoDRI™ ^{5,6} herbicide • Tandem™ herbicide • Traxos® • Trondus® • Varro®



**DOWNLOAD THE 2025 FIELD GUIDE APP FOR
PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.**

CROP ROTATION

- | | | | |
|-----------------|-----------------|---------------------------------|------------------|
| • Alfalfa | • Durum wheat | • Oats | • Soybean |
| • Barley | • Dry bean | • Oriental mustard | • Spring wheat |
| • Brown mustard | • Faba bean | • Peas (<i>field</i>) | • Sunflower |
| • Canola | • Flax | • Potatoes | • Winter wheat |
| • Corn | • Juncea Canola | (<i>except seed potatoes</i>) | • Yellow mustard |

GRAZING AND PRE-HARVEST INTERVAL

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest forage or cut hay within 30 days after application.

⁵ Consult tank-mix partner label for rate-specific claims

⁶ Agral 90 at 0.25% v/v or Bindem at 60 mL/ac is required