

## IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

**Crops / Situations:** Amenity grassland, non-crop areas  
**Maximum Individual Dose:** 200 mL per 10 litres water or 4 L/ha  
**Maximum number of applications:** 1 per year  
**Maximum Total Dose:** 4.0 litres product per hectare per year  
**Latest time of application** 31<sup>st</sup> August

DO NOT USE between 1<sup>st</sup> September and the end of February.

For applications made between 1<sup>st</sup> March and 31<sup>st</sup> May, 1 application may be made per calendar year.  
For applications made between 1<sup>st</sup> June and 31<sup>st</sup> July, only 1 application may be made in a 2 year period.  
For applications made between 1<sup>st</sup> August and 31<sup>st</sup> August, only 1 application may be made in a 3 year period.

DO NOT USE on land that will be grazed by livestock.

DO NOT USE on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment.

DO NOT USE on airfields.

DO NOT apply spot applications to more than 20% of the area.

Users must have received adequate instruction, training and guidance in the safe use of the product and must take all reasonable precautions to protect the health of human beings, creatures and plants and safeguard the environment.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.

## SAFETY INFORMATION

**GARLON ULTRA®**

**UFI: AN65-D0D0-M00C-NQHC**



### WARNING

**Causes serious eye irritation**

**May cause damage to kidneys through prolonged or repeated exposure**

**Very toxic to aquatic life with long lasting effects**

Wash hands thoroughly after handling.

Wear protective gloves / protective clothing / eye protection / face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing.

If eye irritation persists: get medical advice / attention.

Collect spillage.

Contains Triclopyr. May produce an allergic reaction.

Dispose of contents / container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05396

**CORTEVA**  
agriscience



**Garlon® Ultra**

**HERBICIDE**

© Trademarks of Corteva Agriscience and its affiliated companies.

**GROUP 4 HERBICIDES**

**Product Registration Number: PCS No. 05396**

A soluble concentrate containing 12 g ae/litre aminopyralid and 120 g ae/litre triclopyr.

For the control of certain PERENNIAL, HERBACEOUS and WOODY WEEDS in amenity grassland.

### Authorisation Holder:

**Corteva Agriscience UK Limited**

Unit H4, Building H, Melbourn Science Park, Cambridge Road,  
Melbourn, Cambridgeshire, SG8 6HB, United Kingdom.

**24 Hour Emergency Telephone Number: + 353 818 663 627**

Technical Hotline: +44 (0) 800 689 8899,

e-mail: ukhotline@corteva.com

National Poisons Centre: 01 809 2166

### Marketing Company

**Whelehan Crop Protection**

Suite 12 Bunkilla Plaza, Bracetown Business Park

Clonee, Ireland, D15 WR13

Tel: + 353 (1) 574 2410

## SAFETY PRECAUTIONS

### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

### Environmental protection:

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads.)

### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

### READ DIRECTIONS FOR USE ON ATTACHED LEAFLET

**PROTECT FROM FROST**

**FOR PROFESSIONAL USE ONLY**

**Triple Rinse Containers, Puncture And Invert To Dry At Time Of Use**

zone non imprimable

0297001

2509



Artwork Free area for  
in-house print of  
2D data matrix code

NET CONTENTS : 1 LITRE e

DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

WARNINGS

Take extreme care to avoid drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

**SENSITIVE PLANTS** may be harmed by residues of GARLON ULTRA in soil and treated vegetation. Do not apply GARLON ULTRA on or adjacent to soil which may be used as garden top-soil, potting soil, etc. or to grass which may be cut and used as mulch or for compost for horticultural or garden crops. Do not use cuttings from treated grass for mulching or composting.

Cut treated grass must not be removed from site, or used for animal feed, animal bedding, composting or mulching.

Treated grass must not be used for grazing.

On level ground there is negligible lateral movement but do not apply GARLON ULTRA around desirable trees or shrubs, do not spray under the canopy and 1 metre at least from the trunk of desirable trees or shrubs. Care should be taken on slopes to prevent leaching into areas where desirable shrubs, etc are present.

NOTES

Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass. Therefore do not spray in drought, hot or very cold weather conditions.

Do not use on grass less than one year old.

To allow maximum translocation to the roots do not cut grass for 7 days after application.

Clover will be killed by application of GARLON ULTRA.

Wash equipment thoroughly with water and detergent immediately after use.

Non-printed area on page 2 ONLY  
(on the back of 2D code area printed  
on the front page)

HERBACEOUS WEED CONTROL

BROADCAST APPLICATION

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

| Weeds                      | Rates of use<br>litres/ha | Optimum timing of application                                           |
|----------------------------|---------------------------|-------------------------------------------------------------------------|
| Bramble <sup>1</sup>       | 4.0                       | Treat when the bramble is actively growing but is less than 50 cm high. |
| Common mugwort             | 4.0                       | Treat when the mugwort is actively growing and less than 70 cm high.    |
| Common nettle <sup>1</sup> | 2.0                       | Treat when the nettles are actively growing and less than 70 cm high.   |
| Creeping thistle           | 4.0                       | Treat when the thistle is actively growing and less than 70 cm high.    |
| Hogweed <sup>2</sup>       | 4.0                       | Treat when the hogweed is actively growing and less than 70 cm high.    |
| Rosebay willowherb         | 4.0                       | Treat when the willowherb is actively growing and less than 15 cm high. |

<sup>1</sup> A second application may be required the following year.

<sup>2</sup> Moderately susceptible

WOODY AND HERBACEOUS WEED CONTROL

LOCALISED SPOT APPLICATION

WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

| Weeds             | Rate of use<br>mL/10 litres of water | Optimum timing of application          |
|-------------------|--------------------------------------|----------------------------------------|
| Bramble           | 150                                  | Up to 1 metre high.                    |
| Broom             | 200                                  | Up to 1 metre high.                    |
| Buddleia          | 200                                  | Up to 1 metre high.                    |
| Gorse             | 150                                  | Up to 1 metre high.                    |
| Japanese knotweed | 200                                  | 1 metre high, with good foliage cover. |

## **TIMING OF APPLICATION**

GARLON ULTRA should be applied between March and the end of August. The timing of application of GARLON ULTRA is crucial and for good results GARLON ULTRA must be applied to actively growing weeds.

## **APPLICATION**

### **Broadcast Treatment**

GARLON ULTRA should be applied through a tractor-mounted hydraulic sprayer provided it is in good working order and has been calibrated according to the manufacturers' recommendations.

### **Mixing**

Fill the spray tank half full with water and add the required amount of GARLON ULTRA mixing well. Top up with water and continue agitation until the spray tank is full. Maintain agitation while spraying. Use the spray immediately.

### **Spray Volume**

For overall application GARLON ULTRA should be used in a spray volume between 300L and 600L per hectare to give good coverage of the weeds. Higher water volumes are recommended where weed densities are high and where the grass is dense at the time of application.

### **Spray Quality**

Apply as a MEDIUM quality spray as defined by the BCPC system.

### **Spot Treatment**

For localised treatment using a suitable lance from a knapsack or tractor mounted sprayer, use a solution of 150-200 mL of GARLON ULTRA depending on the target weed, per 10 litres of water. Woody weeds should be thoroughly wetted with the spray solution, but spraying until «run-off» will decrease activity. The use of flood jets is recommended to prevent drift. Care should be taken to avoid local overdosing. Woody weeds should not exceed a 1 metre in height.

## **TRADEMARK ACKNOWLEDGEMENTS**

GARLON ULTRA is a trademark of Corteva Agriscience and its affiliated companies. All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

## **Corteva Agriscience Conditions of Supply**

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**To access the Safety Data Sheet for this product scan the QR code or use the weblink below:**



<https://www.corteva.ie/content/dam/dpagco/corteva/eu/ie/en/files/sds/GARLON-ULTRA-IRL-SDS.pdf>

Alternatively contact your supplier

**THIS PAGE HAS BEEN LEFT  
BLANK INTENTIONALLY**

## IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

**Crops / Situations:** Amenity grassland, non-crop areas  
**Maximum Individual Dose:** 200 mL per 10 litres water or 4 L/ha  
**Maximum number of applications:** 1 per year  
**Maximum Total Dose:** 4.0 litres product per hectare per year  
**Latest time of application** 31<sup>st</sup> August

DO NOT USE between 1<sup>st</sup> September and the end of February.

For applications made between 1<sup>st</sup> March and 31<sup>st</sup> May, 1 application may be made per calendar year.

For applications made between 1<sup>st</sup> June and 31<sup>st</sup> July, only 1 application may be made in a 2 year period.

For applications made between 1<sup>st</sup> August and 31<sup>st</sup> August, only 1 application may be made in a 3 year period.

DO NOT USE on land that will be grazed by livestock.

DO NOT USE on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment.

DO NOT USE on airfields.

DO NOT apply spot applications to more than 20% of the area.

Users must have received adequate instruction, training and guidance in the safe use of the product and must take all reasonable precautions to protect the health of human beings, creatures and plants and safeguard the environment.

**Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.**

## SAFETY INFORMATION

Garlon Ultra®

UFI: AN65-D0D0-M00C-NQHC



### WARNING

**Causes serious eye irritation**

**May cause damage to kidneys through prolonged or repeated exposure**

**Very toxic to aquatic life with long lasting effects**

Wash hands thoroughly after handling.

Wear protective gloves / protective clothing / eye protection / face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing.

If eye irritation persists: get medical advice / attention.

Collect spillage.

Contains Tricopyr. May produce an allergic reaction.

Dispose of contents / container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS No. 05396



# Garlon® Ultra

## HERBICIDE

© Trademarks of Corteva Agriscience and its affiliated companies.

GROUP 4 HERBICIDES

**Product Registration Number: PCS No. 05396**

A soluble concentrate containing 12 g ae/litre aminopyralid and 120 g ae/litre tricopyr.

For the control of certain PERENNIAL, HERBACEOUS and WOODY WEEDS in amenity grassland.

### Authorisation Holder:

**Corteva Agriscience UK Limited**

Unit H4, Building H, Melbourn Science Park, Cambridge Road,  
Melbourn, Cambridgeshire, SG8 6HB, United Kingdom.

**24 Hour Emergency Telephone Number: + 353 818 663 627**

Technical Hotline: +44 (0) 800 689 8899,

e-mail: [ukhotline@corteva.com](mailto:ukhotline@corteva.com)

National Poisons Centre: 01 809 2166

### Marketing Company

**Whelehan Crop Protection**

Suite 12 Bunkilla Plaza, Bracetown Business Park

Clonee, Ireland, D15 WR13

Tel: + 353 (1) 574 2410

## SAFETY PRECAUTIONS

### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

### Environmental protection:

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads.)

### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

### READ DIRECTIONS FOR USE ON ATTACHED LEAFLET

**PROTECT FROM FROST**

**FOR PROFESSIONAL USE ONLY**

**Triple Rinse Containers, Puncture And Invert To Dry At Time Of Use**

NET CONTENTS : 1 LITRE e