

## PRINCIPAL FORTE TANK-MIX COMPATIBILITY IN FORAGE MAIZE

**Must be applied with an 95.0 % w/w oil (rapeseed triglycerides) adjuvant i.e. Codacide or Vivolt**

**PLEASE ENSURE YOU READ THE NOTES SECTION BELOW**  
This version supersedes all previous editions

|                                                             |                                                                         |                                                         |
|-------------------------------------------------------------|-------------------------------------------------------------------------|---------------------------------------------------------|
| Normal = Physically Compatible                              | <i>Italics = Physically Compatible with Continuous Agitation</i>        | Red on Blue = Physically Incompatible                   |
| <b>BOLD CAPITALS</b> = Physically & Biologically Compatible | <b>Bold Italic = Biologically compatible with Continuous Agitation</b>  | <b>RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE</b> |
|                                                             | Black on Orange = Do not mix as bacterial survival is severely impacted |                                                         |

### ADJUVANTS

|                     |             |  |  |  |
|---------------------|-------------|--|--|--|
| CODACIDE (1.0 l/ha) | Vivolt 0.2% |  |  |  |
|---------------------|-------------|--|--|--|

### BIOSTIMULANTS

|                   |                 |  |  |  |
|-------------------|-----------------|--|--|--|
| Blue N + Codacide | Blue N + Vivolt |  |  |  |
|-------------------|-----------------|--|--|--|

### HERBICIDES

|                               |                                    |                                                     |  |
|-------------------------------|------------------------------------|-----------------------------------------------------|--|
| <i>Botiga</i>                 | <i>Dual Gold + Codacide/Vivolt</i> | <i>Shield Pro + Starane Hi-Load HL + Codacide *</i> |  |
| <i>Barracuda*</i>             | <i>Gyo + Codacide/Vivolt</i>       | <i>Starane Hi-Load HL + Codacide *</i>              |  |
| <i>CALLISTO*</i>              | <i>Laudis OD + Codacide/Vivolt</i> | <i>Wing-P + Codacide/Vivolt</i>                     |  |
| <i>Diva + Codacide/Vivolt</i> | <i>Shield Pro + Codacide</i>       |                                                     |  |

### TRACE ELEMENTS - tested in mix with Codacide (1.0 l/ha OR Vivolt 0.2%)

|                       |                       |                     |  |
|-----------------------|-----------------------|---------------------|--|
| <i>Amazinc</i>        | <i>Foliar Potash</i>  | <i>Mancuflo</i>     |  |
| <i>Biotrac</i>        | <i>Glytrei MnP</i>    | <i>Mantrac Pro</i>  |  |
| <i>Bortrac</i>        | <i>Gramitrel</i>      | <i>Molytrac</i>     |  |
| <i>Brassitrel Pro</i> | <i>Grow-Pro</i>       | <i>Phosamco Bio</i> |  |
| <i>Canola Bio</i>     | <i>Kinsidro Grow+</i> | <i>Safe N</i>       |  |
| <i>Captrel</i>        | <i>Magflo 300</i>     | <i>Seniphos</i>     |  |
| <i>Croplift Pro</i>   | <i>Magphos</i>        | <i>Stopit</i>       |  |
| <i>Equilibrium</i>    | <i>Maize Boost</i>    | <i>Thiotrac</i>     |  |
| <i>Ferleaf</i>        | <i>Mancozin</i>       | <i>Zintrac</i>      |  |

### COMBINATION MIXES

#### HERBICIDES + TRACE ELEMENTS

|                                   |                                |                                     |  |
|-----------------------------------|--------------------------------|-------------------------------------|--|
| <i>Callisto #+ Kinsidro Grow+</i> | <i>CALLISTO + MAIZE BOOST*</i> | <i>Starane Hi-Load HL + Calfite</i> |  |
|-----------------------------------|--------------------------------|-------------------------------------|--|

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower's risk.

**Important Note:** Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

#### Notes:

\*Do not use with an adjuvant, due to partner label restrictions. Requires the addition of an anti-foam agent.

\*Do not apply if the crop is under stress.

#### MIXING INSTRUCTIONS.

Half fill the spray tank with clean water and start the agitation.

Pour in the required amount of PRINCIPAL FORTE.

Rinse any empty containers thoroughly and add rinsings to the spray tank.

Where tank-mixes are used, and unless directed otherwise, the preferred order of addition of products to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product should be added separately to the half-full sprayer and be fully dispersed before the addition of the next product.

Add the remainder of the water.

Finally, add the required amount of adjuvant and continue agitation.

Maintain agitation until spraying is completed.

Spray solution MUST be used on the day of mixing and must NOT be left overnight.

Follow manufacturer recommendations regarding correct use of the induction bowl.

#### Please note:

Direct contact of WGs with non-diluted tank-mix partners must be avoided.

The above supersedes the instruction on the Codacide label and so negates the need for pre-mixing.

#### KEY CHANGES

Starane Hi-Load HL now approved for use with Principal Forte.; 21May 25 - addition of Kinsidro Grow+ and in mix with Callisto

**USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.**  
**FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL**

Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB.

All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

Principal Forte contains dicamba + rimsulfuron + thifensulfuron

Hotline: 0800 689 8899 | [ukhotline@corteva.com](mailto:ukhotline@corteva.com) | [www.corteva.co.uk](http://www.corteva.co.uk)