

Volunteer Tame Oat Control.

Volunteer oats are extremely competitive in winter wheat and should be controlled early.


Volunteer oats can be more difficult to manage than wild oats and there's a limited choice of products available that effectively control them.

If an oat crop was grown last season, then check fields in autumn for oat volunteers. If they are present, then deal with them early to reduce any impact on yield.

Control of oats with Broadway® Star

Irrespective of variety (winter/spring) or type (wild or volunteers) control with all ALS inhibitor chemistry control declines beyond GS30.

- Optimum timing for control is prior to tillering, mid-tillering at latest.
- Broadway Star + adjuvant will control wild oats up to GS31.
- A Corteva Agriscience trial from 2021/22 showed that autumn was the optimum timing for using Broadway Star + adjuvant to control the following tame oat varieties – Fusion, Elyann, Isabel, Balado, Mascani, Lineout, Southwark, Canyon and Dalguise.
- In the same trial, spring applications resulted in lower control due to less than ideal weather conditions at the time of application, however, spring applications of Broadway Star still gave acceptable levels of control of Fusion, Elyann and Isabel volunteer oats.



Visit us at [corteva.co.uk](https://www.corteva.co.uk)

USE PLANT PROTECTION PRODUCTS SAFELY. Always read the label and product information before use.
 For further information including warning phrases and symbols refer to label.
 Corteva Agriscience, CPC2 Capital Park, Fulbourn, Cambridge CB21 5XE. Tel: 01462 457272.
 ®, ™ Trademarks of Corteva Agriscience and its affiliated companies.
 Technical Hotline: 0800 689 8899. E-mail: ukhotline@corteva.co.uk
 All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
 Broadway Star® contains pyroxulam and florasulam.
 October 2022